

Comments and Addenda

The *Comments and Addenda* section is for short communications which are not of such urgency as to justify publication in *Physical Review Letters* and are not appropriate for regular *Articles*. It includes only the following types of communications: (1) comments on papers previously published in *The Physical Review* or *Physical Review Letters*; (2) addenda to papers previously published in *The Physical Review* or *Physical Review Letters*, in which the additional information can be presented without the need for writing a complete article. Manuscripts intended for this section should be accompanied by a brief abstract for information-retrieval purposes. Accepted manuscripts will follow the same publication schedule as articles in this journal, and galley proofs will be sent to authors.

Revised value for the $\pi \rightarrow e \nu$ branching ratio*

Douglas Bryman and Charles Picciotto

Tri-University Meson Facility and Department of Physics, University of Victoria, Victoria, British Columbia, Canada

(Received 30 September 1974)

Using the current value for the pion lifetime, we have recomputed the branching ratio for $\pi \rightarrow e \nu$ decay obtained from the 1963 experiment of Di Capua *et al.* and find $\Gamma(\pi \rightarrow e \nu + \pi \rightarrow e \nu \gamma) / \Gamma(\pi \rightarrow \mu \nu) = (1.274 \pm 0.024) \times 10^{-4}$, a change of 2%. The theoretically predicted value lies within two standard deviations, compared to the $\frac{1}{2}$ standard deviation reported previously.

The branching ratio measurement for the electronic mode of pion decay is one of the most sensitive tests of the principle of electron-muon universality. The most accurate experiment, performed by Di Capua *et al.*,¹ used a NaI crystal to detect the electrons as well as the photons emitted in internal bremsstrahlung. The measured ratio is

$$\frac{\Gamma(\pi \rightarrow e \nu_e) + \Gamma(\pi \rightarrow e \nu_e \gamma)}{\Gamma(\pi \rightarrow \mu \nu_\mu)}$$

The experimental setup made the measured ratio dependent on the pion decay rate λ_π , such that

$$R_{\text{exp}} = \frac{\lambda_\mu}{\lambda_\pi - \lambda_\mu} \frac{N_{\pi e}}{D e^{\lambda_\mu t_s} - N_{\mu e}} (1 - e^{-(\lambda_\pi - \lambda_\mu) t_s}), \quad (1)$$

where λ_μ is the muon decay rate, D , $N_{\pi e}$, $N_{\mu e}$ are experimentally measured electron yields, and t_s is the separation between two experimental time bins. The measured ratio obtained with the known pion lifetime 25.5 ± 0.3 nsec in 1963 (see Ref. 2) was $R_{\text{exp}} = (1.247 \pm 0.028) \times 10^{-4}$. The theoretically

predicted ratio, taking into account radiative corrections,³ was found to be 1.232×10^{-4} . This agreement is considered a great success of the $V-A$ theory and a confirmation of the hypothesis of electron-muon universality⁴ to better than 1%.

We have recomputed the experimental branching ratio in Eq. (1), using the current value⁵ of the pion lifetime, 26.030 ± 0.023 nsec, to obtain $R_{\text{exp}} = (1.274 \pm 0.024) \times 10^{-4}$, and also the theoretical result, with radiative corrections, using the current value⁵ of the pion mass, to obtain $R_{\text{theo}} = 1.233 \times 10^{-4}$. The theoretical value is within 2 standard deviations of the experimental one, compared to the $\frac{1}{2}$ standard deviation reported previously.

The above theoretical result uses the same ultraviolet cutoff parameter Λ for both the $\pi \rightarrow e \nu$ and $\pi \rightarrow \mu \nu$ virtual-photon corrections. It is interesting to note that if we take the ratio of the values of Λ for the two processes to be in the ratio of the masses of the electron and muon, $\Lambda_e / \Lambda_\mu \simeq m_e / m_\mu$, then the theoretical result becomes $R_{\text{theo}} = 1.258 \times 10^{-4}$, in better agreement with the experiment.

*Work supported in part by the National Research Council of Canada under Grant No. A5926.

¹E. Di Capua *et al.*, Phys. Rev. **133**, B1333 (1964).

²J. Ashkin *et al.*, Nuovo Cimento **16**, 490 (1960).

³T. Kinoshita, Phys. Rev. Lett. **2**, 477 (1959).

⁴See, for example, R. E. Marshak, Riazzudin, and C. P. Ryan, *Theory of Weak Interactions in Particle Physics* (Wiley-Interscience, New York, 1969).

⁵V. Chaloupka *et al.*, Phys. Lett. **50B**, 1 (1974).